NCERT Solutions Class 12 Maths Chapter 10 Vector Algebra

Question 1:

Represent graphically a displacement of 40km, 30° east of north.

Solution:



 \overrightarrow{OP} represents the displacement of 40km, 30° north-east.

Question 2:

Classify the following measures as scalars and vectors.

(i) 10 kg	(ii) 2 meters north-east	(iii) 40°
(iv) 40 watt	(v) 10^{-19} coulomb	(vi) $20m/s^2$

Solution:

- (i) 10kg is a scalar.
- (ii) 2 meters north-west is a vector.
- (iii) 40° is a scalar.
- (iv) 40 watts is a scalar.
- (v) 10^{-19} Coulomb is a scalar.
- (vi) $20m/s^2$ is a vector

Question 3:

Classify the following as scalar and vector quantities.

(i) time period(ii) distance(iii) force(iv) velocity(v) work done.

Solution:

- (i) Time period is a scalar.
- (ii) Distance is a scalar.

- (iii) Force is a vector.
- (iv) Velocity is a vector.
- (v) Work done is a scalar.

Question 4:

In figure, identify the following vectors.



- Vectors \vec{a} and \vec{d} are coinitial. (i)
- Vectors \vec{b} and \vec{d} are equal. (ii)
- (iii) Vectors a and c are collinear but not equal.

Question 5:

Answer the following as true or false.

- a and -a are collinear. (i)
- Two collinear vectors are always equal in magnitude. (ii)
- Two vectors having same magnitude are collinear. (iii)
- Two collinear vectors having the same magnitude are equal. (iv)

Solution:

- (i) True.
- (ii) False.
- (iii) False.
- (iv) False

EXERCISE 10.2

Question 1:

Compute the magnitude of the following vectors:

$$\vec{a} = \hat{i} + \hat{j} + \hat{k}; \quad \vec{b} = \hat{2}\hat{i} - \hat{7}\hat{j} - \hat{3}\hat{k}; \qquad \vec{c} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} - \frac{1}{\sqrt{3}}\hat{k}$$

Solution:

$$\begin{vmatrix} \vec{a} \end{vmatrix} = \sqrt{(1)^2 + (1)^2 + (1)^2} = \sqrt{3} \\ |\vec{b}| = \sqrt{(2)^2 + (-7)^2 + (-3)^2} = \sqrt{4 + 49 + 9} = \sqrt{62} \\ |\vec{c}| = \sqrt{\left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{1}{\sqrt{3}}\right)^2 + \left(-\frac{1}{\sqrt{3}}\right)^2} = \sqrt{\frac{1}{3} + \frac{1}{3} + \frac{1}{3}} = \frac{1}{\sqrt{3}}$$

Question 2:

Write two different vectors having same magnitude.

Solution:

Let
$$\vec{a} = (i - 2j + 3k)$$
 and $\vec{b} = (2i + j - 3k)$
 $|\vec{a}| = \sqrt{1^2 + (-2)^2 + 3^2} = \sqrt{1 + 4 + 9} = \sqrt{14}$
 $|\vec{b}| = \sqrt{2^2 + 1^2 + (-3)^2} = \sqrt{4 + 1 + 9} = \sqrt{14}$

But
$$a \neq b$$

Question 3:

Write two different vectors having same direction.

Solution: Let $\overrightarrow{p} = (\overrightarrow{i} + \overrightarrow{j} + \overrightarrow{k})$ and $\overrightarrow{q} = (2\overrightarrow{i} + 2\overrightarrow{j} + 2\overrightarrow{k})$

The DCs of $\stackrel{\Box}{p}$ are

$$l = \frac{1}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{1}{\sqrt{3}}$$
$$m = \frac{1}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{1}{\sqrt{3}}$$
$$n = \frac{1}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{1}{\sqrt{3}}$$

The DCs of q are

$$l = \frac{2}{\sqrt{2^2 + 2^2 + 2^2}} = \frac{2}{2\sqrt{3}}$$
$$m = \frac{2}{\sqrt{2^2 + 2^2 + 2^2}} = \frac{2}{2\sqrt{3}}$$
$$n = \frac{2}{\sqrt{2^2 + 2^2 + 2^2}} = \frac{2}{2\sqrt{3}}$$

But $p \neq q$

Question 4:

Find the values of x and y so that the vectors 2i+3j and xi+yj are equal.

Solution:

It is given that the vectors 2i+3j and xi+yj are equal.

Therefore,

$$2i+3j = xi + yj$$

On comparing the components of both sides

$$\Rightarrow x = 2$$
$$\Rightarrow y = 3$$

Question 5:

Find the scalar and vector components of the vector with initial point (2,1) and terminal point (-5,7).

Solution:

Let the points be P(2,1) and Q(-5,7)

$$\frac{H}{PQ} = (-5-2)i + (7-1)j$$
$$= -7i + 6j$$

So, the scalar components are -7 and 6, and the vector components are -7i and 6j.

Question 6:

Find the sum of the vectors $\vec{a} = i - 2j + k$, $\vec{b} = -2i - 4j + 5k$ and $\vec{c} = i - 6j + 7k$

Solution:

The given vectors are $\vec{a} = i - 2j + k$, $\vec{b} = -2i - 4j + 5k$ and $\vec{c} = i - 6j + 7k$.

Therefore,

$$\vec{a} + \vec{b} + \vec{c} = (1 - 2 + 1)\hat{i} + (-2 + 4 - 6)\hat{j} + (1 + 5 - 7)\hat{k}$$
$$= \hat{0}\hat{i} - \hat{4}\hat{j} - \hat{k}$$
$$= -\hat{4}\hat{j} - \hat{k}$$

Question 7:

Find the unit vector in the direction of the vector $\vec{a} = i + j + 2k$.

Solution:

We have a = i + j + 2k

Hence,

$$\vec{a} = \sqrt{1^2 + 1^2 + 2^2} = \sqrt{1 + 1 + 4} = \sqrt{6}$$

Therefore,

$$a = \frac{\overrightarrow{a}}{|a|} = \frac{\overrightarrow{i+j+2k}}{\sqrt{6}}$$
$$= \frac{1}{\sqrt{6}}i + \frac{1}{\sqrt{6}}j + \frac{2}{\sqrt{6}}k$$

Question 8:

Find the unit vector in the direction of vector $\stackrel{\text{WD}}{PQ}$, where *P* and *Q* are the points (1,2,3) and (4,5,6) respectively.

Solution:

We have the given points P(1,2,3) and Q(4,5,6)

Hence,

$$\begin{array}{l} \overrightarrow{PQ} = (4-1)i + (5-2)j + (6-3)k \\ = 3i + 3j + 3k \\ |\overrightarrow{PQ}| = \sqrt{3^2 + 3^2 + 3^2} \\ = \sqrt{9 + 9 + 9} \\ = \sqrt{27} \\ = 3\sqrt{3} \end{array}$$

So, unit vector is

$$\frac{PQ}{PQ} = \frac{9i + 3j + 3k}{3\sqrt{3}}$$
$$= \frac{1}{\sqrt{3}}i + \frac{1}{\sqrt{3}}j + \frac{1}{\sqrt{3}}k$$

Question 9:

For given vectors, $\vec{a} = 2i - j + 2k$ and $\vec{b} = -i + j - k$, find the unit vector in the direction of the vector $\vec{a} + \vec{b}$.

Solution:

The given vectors are $\vec{a} = 2i - j + 2k$ and $\vec{b} = -i + j - k$

Therefore,

$$\vec{a} + \vec{b} = (2-1)\hat{i} + (-1+1)\hat{j} + (2-1)\hat{k}$$
$$= \hat{1}\hat{i} + \hat{0}\hat{j} + \hat{1}\hat{k}$$
$$= \hat{i} + \hat{k}$$
$$|\vec{a} + \vec{b}| = \sqrt{1^2 + 1^2} = \sqrt{2}$$

Thus, unit vector is

$$\vec{a+b} = \vec{i+k}$$
$$= \vec{1}\sqrt{2}$$
$$= \frac{1}{\sqrt{2}}i + \frac{1}{\sqrt{2}}k$$

Question 10:

Find a vector in the direction of vector $\hat{5}i - j + \hat{2}k$ which has magnitude 8 units.

Solution:

Let $\vec{a} = \hat{5}i - \hat{j} + \hat{2}k$

Hence,

$$\begin{vmatrix} \vec{a} \\ = \sqrt{5^2 + (-1)^2 + (2)^2} \\ = \sqrt{25 + 1 + 4} \\ = \sqrt{30} \end{vmatrix}$$

Therefore,

$$a = \frac{\overrightarrow{a}}{\left|a\right|} = \frac{\widehat{5i - j + 2k}}{\sqrt{30}}$$

Thus, a vector parallel to $\hat{5i-j+2k}$ with magnitude 8 units is

$$\hat{8}a = 8 \left(\frac{\hat{5}i - j + \hat{2}k}{\sqrt{30}} \right) \\ = \frac{40^{\circ}}{\sqrt{30}}i - \frac{8^{\circ}}{\sqrt{30}}j + \frac{16^{\circ}}{\sqrt{30}}k$$

Question 11:

Show that the vectors $\hat{2}i - \hat{3}j + \hat{4}k$ and $\hat{-4}i + \hat{6}j - \hat{8}k$ are collinear.

Solution:

We have $\vec{a} = \hat{2}i - \hat{3}j + \hat{4}k$ and $\vec{b} = -\hat{4}i + \hat{6}j - \hat{8}k$ Now, $\vec{b} = -\hat{4}i + \hat{6}j - \hat{8}k$

$$p = -4i + 6j - 8k$$
$$= -2(\hat{2}i - \hat{3}j + \hat{4}k)$$
$$= -2a$$

Since, $\vec{b} = \lambda \vec{a}$

Therefore, $\lambda = -2$

So, the vectors are collinear.

Question 12:

Find the direction cosines of the vector $i + \hat{2}j + \hat{3}k$

Solution:

Let $\vec{a} = i + 2j + 3k$

Therefore,

$$\left| \overrightarrow{a} \right| = \sqrt{1^2 + 2^2 + 3^2}$$
$$= \sqrt{1 + 4 + 9}$$
$$= \sqrt{14}$$

Thus, the DCs of \vec{a} are $\left(\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

Question 13:

Find the direction of the cosines of the vectors joining the points A(1,2,-3) and B(-1,-2,1) directions from A to B.

Solution:

The given points are A(1, 2, -3) and B(-1, -2, 1).

Therefore,

$$\begin{array}{l} \begin{array}{l} \blacksquare B \\ \hline AB \\ \hline AB \\ \hline AB \\ \hline AB \\ \hline \hline (-2)^2 + (-4)^2 + 4^2 \\ \hline AB \\ \hline \hline (-2)^2 + (-4)^2 + 4^2 \\ \hline = \sqrt{4 + 16 + 16} \\ \hline = \sqrt{36} \\ \hline = 6 \end{array}$$

Thus, the DCs of
$$AB' \text{ are } \left(-\frac{2}{6}, -\frac{4}{6}, \frac{4}{6}\right) = \left(-\frac{1}{3}, -\frac{2}{3}, \frac{2}{3}\right)$$

Question 14:

Show that the vector i + j + k is equally inclined to the axis OX, OY and OZ.

Solution:

Let $\vec{a} = i + j + k$

Therefore,

$$\left| \overrightarrow{a} \right| = \sqrt{1^2 + 1^2 + 1^2}$$
$$= \sqrt{3}$$

Thus, the DCs of a are $\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$

Now, let α , β and γ be the angles formed by a with the positive directions of x, y and z axes respectively

Then,

$$\cos \alpha = \frac{1}{\sqrt{3}}, \cos \beta = \frac{1}{\sqrt{3}}, \cos \gamma = \frac{1}{\sqrt{3}}$$

Hence, the vector is equally inclined to OX, OY and OZ.

Question 15:

Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are i+2j-k and -i+j+k respectively, in the ratio 2:1.

- (i) Internally
- (ii) Externally

Solution:

Position vectors of P and Q are given as:

$$\overrightarrow{OP} = i + 2j - k$$
 and $\overrightarrow{OQ} = -i + j + k$

(i) The position vector of R which divides the line joining two points P and Q internally in the ratio 2:1 is

$$\overrightarrow{OR} = \frac{2(\hat{-i}+\hat{j}+\hat{k})+1(\hat{i}+\hat{2}\hat{j}-\hat{k})}{2+1} \\
 = \frac{(\hat{-2i}+\hat{2}\hat{j}+\hat{2}\hat{k})+(\hat{i}+\hat{2}\hat{j}-\hat{k})}{3} \\
 = \frac{\hat{-i}+\hat{4}\hat{j}+\hat{k}}{3} \\
 = -\frac{\hat{r}}{3}\hat{i}+\frac{\hat{4}}{3}\hat{j}+\frac{\hat{r}}{3}\hat{k}$$

(ii) The position vector of R which divides the line joining two points P and Q externally in the ratio 2:1 is

$$\overrightarrow{OR} = \frac{2(\hat{-i}+\hat{j}+\hat{k})-1(\hat{i}+\hat{2}\hat{j}-\hat{k})}{2-1} \\ = \frac{(\hat{-2}\hat{i}+\hat{2}\hat{j}+\hat{2}\hat{k})-(\hat{i}+\hat{2}\hat{j}-\hat{k})}{1} \\ = -\hat{3}\hat{i}+\hat{3}\hat{k}$$

Question 16:

Find the position vector of the mid-point of the vector joining the points P(2,3,4) and Q(4,1,-2).

Solution:

The position vector of the mid-point R is

Question 17:

Show that the points A,B and C with position vectors, $\vec{a} = \hat{3}i - \hat{4}j - \hat{4}k$, $\vec{b} = \hat{2}i - \hat{j} + \hat{k}$ and $\vec{c} = i - \hat{3}j - \hat{5}k$, respectively form the vertices of a right angled triangle.

Solution:

Position vectors of points A, B, and C are respectively given as:

$$\vec{a} = \hat{3}i - \hat{4}j - \hat{4}k$$
, $\vec{b} = \hat{2}i - \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - \hat{3}j - \hat{5}k$

Therefore,

Now,

$$\begin{aligned} \underbrace{\square}_{AB}^{2} &= (-1)^{2} + 3^{2} + 5^{2} = 1 + 9 + 25 = 35 \\ \underbrace{\square}_{BC}^{2} &= (-1)^{2} + (-2)^{2} + (-6)^{2} = 1 + 4 + 36 = 41 \\ \underbrace{\square}_{CA}^{2} &= 2^{2} + (-1)^{2} + 1^{2} = 4 + 1 + 1 = 6 \end{aligned}$$

Also,

$$\begin{aligned} \overrightarrow{AB}^2 + |\overrightarrow{CA}|^2 &= 35 + 6 \\ &= 41 \\ &= |\overrightarrow{BC}|^2 \end{aligned}$$

Thus, ABC is a right-angled triangle.

Question 18:

In triangle ABC which of the following is not true.



- (A) $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0$

- (A) AB + BC + CA = 0(B) AB + BC AC = 0(C) AB + BC CA = 0(D) AB CB + CA = 0

Solution:



On applying the triangle law of addition in the given triangle, we have:

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC} \qquad \dots(1)$$

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} = -\overrightarrow{CA}$$

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} = -\overrightarrow{CA}$$

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0 \qquad \dots(2)$$

Hence, the equation given in option A is true.

Now, from equation (2)

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0$$

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} - \overrightarrow{AC} = 0$$

Hence, the equation given in option B is true.

Also,

$$\Rightarrow \overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA} = 0$$

$$\implies \overrightarrow{AB} - \overrightarrow{CB} + \overrightarrow{CA} = 0$$

Hence, the equation given in option D is true

Now, consider the equation given in option C,

$$\begin{array}{c} \blacksquare P \\ AB + BC - CA = 0 \\ \blacksquare P \\ \blacksquare P \\ \Rightarrow AB + BC = CA \\ \end{array}$$
...(3)

From equations (1) and (2)

$$\Rightarrow \overrightarrow{AC} = \overrightarrow{CA}$$

$$\Rightarrow \overrightarrow{AC} = -\overrightarrow{AC}$$

$$\Rightarrow \overrightarrow{AC} = -\overrightarrow{AC}$$

$$\Rightarrow \overrightarrow{AC} + \overrightarrow{AC} = 0$$

$$\Rightarrow 2\overrightarrow{AC} = 0$$

$$\Rightarrow \overrightarrow{AC} = 0$$

Which is not true. So, the equation given in option C is incorrect.

Thus, the correct option is C.

Question 19:

If a and b are two collinear vectors, then which of the following are incorrect?

- (A) $b = \lambda a$, for some scalar λ
- (B) $a = \pm b$
- (c) the respective components of \vec{a} and \vec{b} are proportional.
- (D) both the vectors \vec{a} and \vec{b} have same direction, but different magnitudes

Solution:

If \vec{a} and \vec{b} are collinear vectors, they are parallel.

Therefore, for some scalar λ

$$b = \lambda a$$

If $\lambda = \pm 1$, then $a = \pm b$

If
$$\vec{a} = \hat{a_1}\hat{i} + \hat{a_2}\hat{j} + \hat{a_3}\hat{k}$$
 and $\vec{b} = \hat{b_1}\hat{i} + \hat{b_2}\hat{j} + \hat{b_3}\hat{k}$

Then,

$$\Rightarrow \hat{b} = \lambda \hat{a}$$

$$\Rightarrow \hat{b_1}i + \hat{b_2}j + \hat{b_3}k = \lambda \left(\hat{a_1}i + \hat{a_2}j + \hat{a_3}k\right)$$

$$\Rightarrow \hat{b_1}i + \hat{b_2}j + \hat{b_3}k = (\lambda \hat{a_1})i + (\lambda \hat{a_2})j + (\lambda \hat{a_3})k$$

Comparing the components of both the sides

$$\Rightarrow b_1 = \lambda a_1$$

$$\Rightarrow b_2 = \lambda a_2$$

$$\Rightarrow b_3 = \lambda a_3$$

Therefore,

$$\frac{b_1}{a_1} = \frac{b_2}{a_2} = \frac{b_3}{a_3} = \lambda$$

Thus, the respective components of \vec{a} and \vec{b} are proportional.

However, a and \overline{b} may have different directions.

Hence, that statement given in D is incorrect.

Thus, the correct option is D.

EXERCISE 10.3

Question 1:

Find the angle between two vectors \vec{a} and \vec{b} with magnitude $\sqrt{3}$ and 2, respectively have $\vec{a}.\vec{b} = \sqrt{6}$

Solution:

It is given that

$$\begin{vmatrix} \overrightarrow{a} \\ = \sqrt{3} \\ |\overrightarrow{b}| = 2 \\ \overrightarrow{a.b} = \sqrt{6} \end{vmatrix}$$

Therefore,

$$\Rightarrow \sqrt{6} = \sqrt{3} \times 2\cos\theta$$
$$\Rightarrow \cos\theta = \frac{\sqrt{6}}{\sqrt{3} \times 2}$$
$$\Rightarrow \cos\theta = \frac{1}{\sqrt{2}}$$
$$\Rightarrow \theta = \frac{\pi}{4}$$

Question 2:

Find the angle between the vectors i - 2j + 3k and 3i - 2j + k.

Solution:

Let $\vec{a} = i - 2j + 3k$ and $\vec{b} = 3i - 2j + k$.

Hence,

$$\begin{vmatrix} \vec{a} \end{vmatrix} = \sqrt{1^2 + (-2)^2 + 3^2} = \sqrt{1 + 4 + 9} = \sqrt{14} \\ |\vec{b}| = \sqrt{3^2 + (-2)^2 + 1^2} = \sqrt{9 + 4 + 1} = \sqrt{14} \\ \vec{a}.\vec{b} = (i - 2j + 3k)(3i - 2j + k) \\ = 1 \times 3 + (-2)(-2) + 3 \times 1 \\ = 3 + 4 + 3 \\ = 10 \end{vmatrix}$$

Therefore,

$$\Rightarrow 10 = \sqrt{14}\sqrt{14}\cos\theta$$
$$\Rightarrow \cos\theta = \frac{10}{14}$$
$$\Rightarrow \theta = \cos^{-1}\left(\frac{5}{7}\right)$$

Question 3:

Find the projection of the vector $\hat{i-j}$ on the vector $\hat{i+j}$

Solution:

Let $\vec{a} = i - j$ and $\vec{b} = i + j$

Projection of \vec{a} on \vec{b} is

$$\frac{1}{|b|} (\overrightarrow{a.b}) = \frac{1}{\sqrt{1+1}} \{ (1)(1) + (-1)1 \}$$
$$= \frac{1}{\sqrt{2}} (1-1)$$
$$= 0$$

Question 4:

Find the projection of vector i+3j+7k on the vector $\hat{7}i-j+\hat{8}k$

Solution:

Let $\vec{a} = i + 3j + 7k$ and $\vec{b} = 7i - j + 8k$

Projection of \vec{a} on \vec{b} is

$$\frac{1}{|b|} (\overrightarrow{a.b}) = \frac{1}{\sqrt{7^2 + (-1)^2 + 8^2}} \{ (1)(7) + 3(-1) + 7(8) \}$$
$$= \frac{1}{\sqrt{49 + 1 + 64}} (7 - 3 + 56)$$
$$= \frac{60}{\sqrt{114}}$$

Question 5:

Show that each of the given three vectors is a unit vector which are manually perpendicular to each other.

$$\frac{1}{7}(\hat{2}i+\hat{3}j+\hat{6}k),\frac{1}{7}(\hat{3}i-\hat{6}j+\hat{2}k),\frac{1}{7}(\hat{6}i+\hat{2}j-\hat{3}k)$$

Solution:

Let

$$\vec{a} = \frac{1}{7} \left(\hat{2}i + \hat{3}j + \hat{6}k \right) = \frac{2}{7}i + \frac{3}{7}j + \frac{6}{7}k$$
$$\vec{b} = \frac{1}{7} \left(\hat{3}i - \hat{6}j + \hat{2}k \right) = \frac{3}{7}i - \frac{6}{7}j + \frac{2}{7}k$$
$$\vec{c} = \frac{1}{7} \left(\hat{6}i + \hat{2}j - \hat{3}k \right) = \frac{6}{7}i + \frac{2}{7}j - \frac{3}{7}k$$

Now,

$$\begin{vmatrix} \vec{a} \end{vmatrix} = \sqrt{\left(\frac{2}{7}\right)^2 + \left(\frac{3}{7}\right)^2 + \left(\frac{6}{7}\right)^2} = \sqrt{\frac{4}{49} + \frac{9}{49} + \frac{36}{49}} = 1$$
$$\begin{vmatrix} \vec{b} \end{vmatrix} = \sqrt{\left(\frac{3}{7}\right)^2 + \left(-\frac{6}{7}\right)^2 + \left(\frac{2}{7}\right)^2} = \sqrt{\frac{9}{49} + \frac{36}{49} + \frac{4}{49}} = 1$$
$$\begin{vmatrix} \vec{c} \end{vmatrix} = \sqrt{\left(\frac{6}{7}\right)^2 + \left(\frac{2}{7}\right)^2 + \left(-\frac{3}{7}\right)^2} = \sqrt{\frac{36}{49} + \frac{4}{49} + \frac{9}{49}} = 1$$

So, each of the vector is a unit vector.

Hence,

$$\overrightarrow{a.b} = \frac{2}{7} \times \frac{3}{7} + \frac{3}{7} \times \left(-\frac{6}{7}\right) + \frac{6}{7} \times \frac{2}{7} = \frac{6}{49} - \frac{18}{49} + \frac{12}{49} = 0$$

$$\overrightarrow{b.c} = \frac{3}{7} \times \frac{6}{7} + \frac{2}{7} \times \left(-\frac{6}{7}\right) + \left(-\frac{3}{7}\right) \times \frac{2}{7} = \frac{18}{49} - \frac{12}{49} - \frac{6}{49} = 0$$

$$\overrightarrow{c.a} = \frac{2}{7} \times \frac{6}{7} + \frac{3}{7} \times \frac{2}{7} + \frac{6}{7} \times \left(-\frac{3}{7}\right) = \frac{12}{49} + \frac{6}{49} + \frac{18}{49} = 0$$

So, the vectors are mutually perpendicular to each other.

Question 6:

Find $|\vec{a}|_{\text{and}} |\vec{b}|_{\text{, if }} (\vec{a} + \vec{b}) (\vec{a} - \vec{b}) = 8$ and $|\vec{a}| = 8|\vec{b}|$

Solution:

It is given that $(\vec{a} + \vec{b})(\vec{a} - \vec{b}) = 8$ and $|\vec{a}| = 8|\vec{b}|$ Therefore,

$$\Rightarrow (\vec{a} + \vec{b})(\vec{a} - \vec{b}) = 8$$

$$\Rightarrow \vec{a} \cdot \vec{a} - \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} - \vec{b} \cdot \vec{b} = 8$$

$$\Rightarrow |\vec{a}|^2 - |\vec{b}|^2 = 8$$

$$\Rightarrow (8|\vec{b}|)^2 - |\vec{b}|^2 = 8$$

$$\Rightarrow 64|\vec{b}|^2 - |\vec{b}|^2 = 8$$

$$\Rightarrow 63|\vec{b}|^2 = 9$$

$$\Rightarrow |\vec{b}|^2 = \frac{8}{63}$$

$$\Rightarrow |\vec{b}| = \sqrt{\frac{8}{63}}$$

$$\Rightarrow |\vec{b}| = \sqrt{\frac{8}{53}}$$

Now,

$$\begin{vmatrix} a \end{vmatrix} = 8 \begin{vmatrix} b \end{vmatrix}$$
$$= \frac{8 \times 2\sqrt{2}}{3\sqrt{7}}$$
$$= \frac{16\sqrt{2}}{3\sqrt{7}}$$

Question 7:

Evaluate the product (3a-5b)(2a+7b)

Solution:

$$(3\overrightarrow{a}-5\overrightarrow{b})(2\overrightarrow{a}+7\overrightarrow{b}) = 3\overrightarrow{a}.2\overrightarrow{a}+3\overrightarrow{a}.7\overrightarrow{b}-5\overrightarrow{b}.2\overrightarrow{a}-5\overrightarrow{b}.7\overrightarrow{b}$$
$$= 6\overrightarrow{a}\overrightarrow{a}+21\overrightarrow{a}\overrightarrow{b}-10\overrightarrow{a}\overrightarrow{b}-35\overrightarrow{b}\overrightarrow{b}$$
$$= 6\left|\overrightarrow{a}\right|^{2}+11\overrightarrow{a}\overrightarrow{b}-35\left|\overrightarrow{b}\right|^{2}$$

Question 8:

Find the magnitude of two vectors \vec{a} and \vec{b} , having the same magnitude and such that angle between them is 60° and their scalar product is $\frac{1}{2}$

Solution:

Let θ be the angle between \vec{a} and \vec{b}

It is given that $|\vec{a}| = |\vec{b}|$, $\vec{a.b} = \frac{1}{2}$ and $\theta = 60^{\circ}$ Therefore,

$$\Rightarrow \frac{1}{2} = |\vec{a}|\vec{b}| \cos 60^{\circ}$$
$$\Rightarrow \frac{1}{2} = |\vec{a}|^{2} \times \frac{1}{2}$$
$$\Rightarrow |\vec{a}|^{2} = 1$$
$$\Rightarrow |\vec{a}| = |\vec{b}| = 1$$

Question 9:

Find $|\vec{x}|$, if for a unit vector \vec{a} , $(\vec{x}-\vec{a}) \cdot (\vec{x}+\vec{a}) = 12$

Solution:

$$\Rightarrow (\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$$

$$\Rightarrow \vec{x} \cdot \vec{x} + \vec{x} \cdot \vec{a} - \vec{a} \cdot \vec{x} - \vec{a} \vec{a} = 12$$

$$\Rightarrow |\vec{x}|^2 - |\vec{a}|^2 = 12$$

$$\Rightarrow |\vec{x}|^2 - 1 = 12$$

$$\Rightarrow |\vec{x}|^2 = 13$$

$$\Rightarrow |\vec{x}| = \sqrt{13}$$

Question 10:

If $\vec{a} = \hat{2}i + \hat{2}j + \hat{3}k$, $\vec{b} = -i + \hat{2}j + \hat{k}$ and $\vec{c} = \hat{3}i + j$ are such that $\vec{a} + \lambda \vec{b}$ is perpendicular to \vec{c} , then find the value of λ .

Solution:

We have $\vec{a} = 2i + 2j + 3k$, $\vec{b} = -i + 2j + k$ and $\vec{c} = 3i + j$ are such that $\vec{a} + \lambda \vec{b}$ is perpendicular to \vec{c}

Then,

$$\vec{a} + \lambda \vec{b} = (\hat{2}i + \hat{2}j + \hat{3}k) + \lambda (\hat{-i} + \hat{2}j + \hat{k})$$
$$= (2 - \lambda)\hat{i} + (2 + 2\lambda)\hat{j} + (3 + \lambda)\hat{k}$$

Now,

$$\Rightarrow (\vec{a} + \lambda \vec{b}) \cdot \vec{c} = 0$$

$$\Rightarrow \left[(2 - \lambda)\hat{i} + (2 + 2\lambda)\hat{j} + (3 + \lambda)\hat{k} \right] \cdot (\hat{3}\hat{i} + \hat{j}) = 0$$

$$\Rightarrow 3(2 - \lambda) + (2 + 2\lambda) + 0(3 + \lambda) = 0$$

$$\Rightarrow 6 - 3\lambda + 2 + 2\lambda = 0$$

$$\Rightarrow -\lambda + 8 = 0$$

$$\Rightarrow \lambda = 8$$

Question 11:

Show that $|\vec{a}|\vec{b}+|\vec{b}|\vec{a}|$ is perpendicular to $|\vec{a}|\vec{b}-|\vec{b}|\vec{a}|$, for any non-zero vectors \vec{a} and \vec{b} .

Solution:

$$(|\vec{a}|\vec{b}+|\vec{b}|\vec{a}).(|\vec{a}|\vec{b}-|\vec{b}|\vec{a}) = |\vec{a}|^2 \vec{b}.\vec{b}-|\vec{a}||\vec{b}|\vec{b}.\vec{a}+|\vec{b}||\vec{a}|\vec{a}.\vec{b}-|\vec{b}|^2 \vec{a}.\vec{a}$$

 $= |\vec{a}|^2 |\vec{b}|^2 - |\vec{b}|^2 |\vec{a}|^2$
 $= 0$

Question 12:

If aa = 0 and ab = 0, then what can be concluded above the vector \vec{b} ?

Solution:

We have a.a = 0 and a.b = 0

Hence,

$$\Rightarrow \left| \overrightarrow{a} \right|^2 = 0$$
$$\Rightarrow \left| \overrightarrow{a} \right| = 0$$

Therefore, \overline{a} is the zero vector

Thus, any vector \vec{b} can satisfy $\vec{a}.\vec{b} = 0$

Question 13:

If a, b, c are unit vectors such that a+b+c=0, find the value of a.b+b.c+c.a.

Solution:

We have a, b, c are unit vectors such that a+b+c=0

Therefore,

$$\begin{vmatrix} \overrightarrow{a} + \overrightarrow{b} + \overrightarrow{c} \end{vmatrix}^{2} = (\overrightarrow{a} + \overrightarrow{b} + \overrightarrow{c}) \cdot (\overrightarrow{a} + \overrightarrow{b} + \overrightarrow{c})$$
$$0 = |\overrightarrow{a}|^{2} + |\overrightarrow{b}|^{2} + |\overrightarrow{c}|^{2} + 2(\overrightarrow{a} \cdot \overrightarrow{b} + \overrightarrow{b} \cdot \overrightarrow{c} + \overrightarrow{c} \cdot \overrightarrow{a})$$
$$0 = 1 + 1 + 1 + 2(\overrightarrow{a} \cdot \overrightarrow{b} + \overrightarrow{b} \cdot \overrightarrow{c} + \overrightarrow{c} \cdot \overrightarrow{a})$$
$$(\overrightarrow{a} \cdot \overrightarrow{b} + \overrightarrow{b} \cdot \overrightarrow{c} + \overrightarrow{c} \cdot \overrightarrow{a}) = \frac{-3}{2}$$

Question 14:

If either vector $\vec{a} = 0$ or $\vec{b} = 0$, then $\vec{a}.\vec{b} = 0$. But the converse need not be true. Justify the answer with an example.

Solution:

Let $\vec{a} = 2i + \hat{4}j + \hat{3}k$ and $\vec{b} = \hat{3}i + \hat{3}j - \hat{6}k$

Therefore,

$$ab = 2(3) + 4(3) + 3(-6)$$

= 6 + 12 - 18
= 0

Now,

$$\begin{vmatrix} a \end{vmatrix} = \sqrt{2^2 + 4^2 + 3^2} = \sqrt{29} \\ \Rightarrow a \neq 0 \\ |\vec{b}| = \sqrt{3^2 + 3^2 + (-6)^2} = \sqrt{54} \\ \Rightarrow \vec{b} \neq 0 \end{vmatrix}$$

So, the converse of the statement need not to be true.

Question 15:

If the vertices A, B, C of a triangle ABC are (1,2,3), (-1,0,0), (0,1,2) respectively, then find $\angle ABC$. $[\angle ABC$ is the angle between the vectors \overrightarrow{BA} and \overrightarrow{BC}]

Solution:

Vertices of the triangle are A(1,2,3), B(-1,0,0) and C(0,1,2).

Hence,

$$BA = \{1 - (1)\}i + (2 - 0)j + (3 - 0)k$$

$$= 2i + 2j + 3k$$

$$BC = \{0 - (-1)\}i + (1 - 0)j + (2 - 0)k$$

$$= i + j + 2k$$

$$BABC = (2i + 2j + 3k)(i + j + 2k)$$

$$= 2 \times 1 + 2 \times 1 + 3 \times 2$$

$$= 2 + 2 + 6$$

$$= 10$$

$$|BA| = \sqrt{2^{2} + 2^{2} + 3^{2}}$$

$$= \sqrt{4 + 4 + 9}$$

$$= \sqrt{17}$$

$$|BC| = \sqrt{1 + 1 + 2^{2}}$$

$$= \sqrt{6}$$

$$BABC = |BA| |BC| \cos(\angle ABC)$$

Therefore,

$$\Rightarrow 10 = \sqrt{17} \times \sqrt{6} \left(\cos \angle ABC \right)$$
$$\Rightarrow \cos \left(\angle ABC \right) = \frac{10}{\sqrt{17} \times \sqrt{6}}$$
$$\Rightarrow \left(\angle ABC \right) = \cos^{-1} \left(\frac{10}{\sqrt{102}} \right)$$

Question 16:

Show that the points A(1,2,7), B(2,6,3) and C(3,10-1) are collinear.

Solution:

The given points are A(1,2,7), B(2,6,3) and C(3,10-1).

Hence,

$$\begin{aligned} \overrightarrow{AB} &= (2-1)i + (6-2)j + (3-7)k = i + 4j - 4k \\ \overrightarrow{BC} &= (3-2)i + (10-6)j + (-1-3)k = i + 4j - 4k \\ \overrightarrow{AC} &= (3-1)i + (10-2)j + (-1-7)k = 2i + 8j - 8k \\ |\overrightarrow{AB}| &= \sqrt{1^2 + 4^2 + (-4)^2} = \sqrt{1 + 16 + 16} = \sqrt{33} \\ |\overrightarrow{BC}| &= \sqrt{1^2 + 4^2 + (-4)^2} = \sqrt{1 + 16 + 16} = \sqrt{33} \\ |\overrightarrow{BC}| &= \sqrt{2^2 + 8^2 + 8^2} = \sqrt{4 + 64 + 64} = 2\sqrt{33} \end{aligned}$$

Therefore,

$$\begin{vmatrix} \mathbf{AB} \\ \mathbf{AB} \end{vmatrix} + \begin{vmatrix} \mathbf{BC} \\ \mathbf{BC} \end{vmatrix} = \sqrt{33} + \sqrt{33}$$
$$= 2\sqrt{33}$$
$$= \begin{vmatrix} \mathbf{AC} \end{vmatrix}$$

Hence, the points are collinear.

Question 17:

Show that the vectors $\hat{2}i - \hat{j} + \hat{k}$, $i - \hat{3}j - \hat{5}k$ and $\hat{3}i - \hat{4}j - \hat{4}k$ form the vertices of a right angled triangle.

Solution:

Let
$$\overrightarrow{OA} = 2i - j + k$$
, $\overrightarrow{OB} = i - 3j - 5k$ and $\overrightarrow{OC} = 3i - 4j - 4k$

Hence,

$$\begin{aligned} \overrightarrow{AB} &= (1-2)i + (-3+1)j + (-5-1)k = -i - 2j - 6k \\ \overrightarrow{BC} &= (3-1)i + (-4+3)j + (-4+5)k = 2i - j + k \\ \overrightarrow{AC} &= (2-3)i + (-1+4)j + (1+4)k = -i + 3j + 5k \\ |\overrightarrow{AC}| &= \sqrt{(-1)^2 + (-2)^2 + (-6)^2} = \sqrt{1+4+36} = \sqrt{41} \\ |\overrightarrow{BC}| &= \sqrt{2^2 + (-1)^2 + 1^2} = \sqrt{4+1+1} = \sqrt{6} \\ |\overrightarrow{AC}| &= \sqrt{(-1)^2 + 3^2 + 5^2} = \sqrt{1+9+25} = \sqrt{35} \end{aligned}$$

Therefore,

$$\begin{vmatrix} \blacksquare \uparrow 1 \\ BC \end{vmatrix}^{2} + \begin{vmatrix} \blacksquare \downarrow \uparrow 2 \\ AC \end{vmatrix}^{2} = 6 + 35$$
$$= 41$$
$$= \begin{vmatrix} \blacksquare \uparrow \blacksquare \end{vmatrix}^{2}$$

Thus, $\triangle ABC$ is a right-angled triangle.

Question 18:

If \vec{a} is a nonzero vector of magnitude 'a' and λ a nonzero scalar, then $\lambda \vec{a}$ is a unit vector if

(A) $\lambda = 1$ (B) $\lambda = -1$ (C) $a = |\lambda|$ (D) $a = \frac{1}{|\lambda|}$

Solution:

 $\Rightarrow \left| \lambda \overrightarrow{a} \right| = 1$ $\Rightarrow \left| \lambda \right| \left| \overrightarrow{a} \right| = 1$ $\Rightarrow \left| \overrightarrow{a} \right| = \frac{1}{\left| \lambda \right|}$ $\Rightarrow a = \frac{1}{\left| \lambda \right|}$

Hence the correct option is D.

EXERCISE 10.4

Question 1: Find $|\vec{a} \times \vec{b}|$, if $\vec{a} = i - \hat{7}j + \hat{7}k$ and $\vec{b} = \hat{3}i - \hat{2}j + \hat{2}k$

Solution: We have, $\vec{a} = i - \hat{7}j + \hat{7}k$ and $\vec{b} = \hat{3}i - \hat{2}j + \hat{2}k$ Hence,

$$\vec{a} \times \vec{b} = \begin{vmatrix} i & j & k \\ 1 & -7 & 7 \\ 3 & -2 & 2 \end{vmatrix}$$
$$= i(-14+14) - j(2-21) + k(-2+21)$$
$$= 19 j + 19k$$

Therefore,

$$\vec{a} \times \vec{b} = \sqrt{(19)^2 + (19)^2}$$
$$= \sqrt{2 \times (19)^2}$$
$$= 19\sqrt{2}$$

Question 2:

Find a unit vector perpendicular to each of the vector $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$, where $\vec{a} = \hat{3}i + \hat{2}j + \hat{2}k$ and $\vec{b} = i + \hat{2}j - \hat{2}k$.

Solution:

We have $\vec{a} = \hat{3}i + \hat{2}j + \hat{2}k$ and $\vec{b} = i + \hat{2}j - \hat{2}k$. Hence,

$$\vec{a} + \vec{b} = 4\vec{i} + 4\vec{j}$$

$$\vec{a} - \vec{b} = 2\vec{i} + 4\vec{k}$$

Therefore,

$$(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b}) = \begin{vmatrix} i & j & k \\ 4 & 4 & 0 \\ 2 & 0 & 4 \end{vmatrix}$$
$$= i(16) - j(16) + k(-8)$$
$$= 16i - 16j - 8k$$
$$|(\vec{a} + \vec{b}) \times (\vec{a} - \vec{b})| = \sqrt{16^2 + (-16)^2 + (-8)^2} = \sqrt{2^2 \times 8^2 + 2^2 \times 8^2 + 8^2}$$
$$= 8\sqrt{2^2 + 2^2 + 1} = 8\sqrt{9}$$
$$= 8 \times 3 = 24$$

So, the unit vector is

$$\pm \frac{\widehat{(a+b)} \times \widehat{(a-b)}}{|\widehat{(a+b)} \times \widehat{(a-b)}|} = \pm \frac{\widehat{16}i - \widehat{16}j - \widehat{8}k}{24}$$
$$= \pm \frac{\widehat{2}i - \widehat{2}j - k}{3}$$
$$= \pm \frac{\widehat{2}i - \widehat{2}j - k}{3}$$

Question 3:

If a unit vector \vec{a} makes an angle $\frac{\pi}{3}$ with i, $\frac{\pi}{4}$ with j and an acute angle θ with k, then find θ and hence, the components of \vec{a} .

Solution:

Let the unit vector $\vec{a} = \hat{a_1}\hat{i} + \hat{a_2}\hat{j} + \hat{a_3}\hat{k}$ Then, $|\vec{a}| = 1$

Now,

$$\cos \frac{\pi}{3} = \frac{a_1}{|a|} \Rightarrow a_1 = \frac{1}{2}$$
$$\cos \frac{\pi}{4} = \frac{a_2}{|a|} \Rightarrow a_2 = \frac{1}{\sqrt{2}}$$
$$\cos \theta = \frac{a_3}{|a|} \Rightarrow a_3 = \cos \theta$$

Therefore,

Hence,

$$\Rightarrow \sqrt{a_1^2 + a_2^2 + a_3^2} = 1$$

$$\Rightarrow \left(\frac{1}{2}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2 + \cos^2 \theta = 1$$

$$\Rightarrow \frac{1}{4} + \frac{1}{2} + \cos^2 \theta = 1$$

$$\Rightarrow \frac{3}{4} + \cos^2 \theta = 1$$

$$\Rightarrow \cos^2 \theta = 1 - \frac{3}{4} = \frac{1}{4}$$

$$\Rightarrow \cos \theta = \frac{1}{2}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

$$a_3 = \cos \frac{\pi}{3} = \frac{1}{2}$$

So, $\theta = \frac{\pi}{3}$ and components of a are $\left(\frac{1}{2}, \frac{1}{\sqrt{2}}, \frac{1}{2}\right)$

Question 4: Show that $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2(\vec{a} \times \vec{b})$

Solution:

$$LHS = (\overrightarrow{a-b}) \times (\overrightarrow{a+b})$$

$$= (\overrightarrow{a-b}) \times \overrightarrow{a+} (\overrightarrow{a-b}) \times \overrightarrow{b}$$

$$= \overrightarrow{a \times a - b \times a + a \times b - b \times b}$$

$$= 0 + a \times b + a \times b - 0$$

$$= 2a \times b$$

$$= RHS$$

Question 5:

Find
$$\lambda$$
 and μ if $(\hat{2}i+\hat{6}j+2\hat{7}k)\times(\hat{i}+\hat{\lambda}j+\hat{\mu}k)=0$

Solution:

We have $(\hat{2}i+\hat{6}j+2\hat{7}k)\times(\hat{i}+\hat{\lambda}j+\hat{\mu}k)=0$ Therefore, $\Rightarrow (\hat{2}i+\hat{6}j+2\hat{7}k)\times(\hat{i}+\hat{\lambda}j+\hat{\mu}k)=0$ $\Rightarrow \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 6 & 27 \\ 1 & \lambda & \mu \end{vmatrix} = \hat{0}i+\hat{0}j+\hat{0}k$

$$\Rightarrow \hat{i}(6\mu - 27\lambda) - \hat{j}(2\mu - 27) + \hat{k}(2\lambda - 6) = \hat{0}\hat{i} + \hat{0}\hat{j} + \hat{0}\hat{k}$$

On comparing the corresponding components, we have:

$$6\mu - 27\lambda = 0$$
$$2\mu - 27 = 0$$
$$2\lambda - 6 = 0$$

Now,

$$2\lambda - 6 = 0 \Longrightarrow \lambda = 3$$
$$2\mu - 27 = 0 \Longrightarrow \mu = \frac{27}{2}$$

Question 6:

Given that a.b=0 and $a \times b = 0$. What can you conclude about a and b?

Solution:

When a.b = 0

Either $|\vec{a}| = 0$ or $|\vec{b}| = 0$ Or $\vec{a} \perp \vec{b}$ (if $|\vec{a}| \neq 0$ and $|\vec{b}| \neq 0$) When $\vec{a} \times \vec{b} = 0$ Either $|\vec{a}| = 0$ or $|\vec{b}| = 0$ Or $\vec{a} \parallel \vec{b}$ (if $|\vec{a}| \neq 0$ and $|\vec{b}| \neq 0$)

Since, \vec{a} and \vec{b} cannot be perpendicular and parallel simultaneously.

So, $\vec{a} = 0$ or $\vec{b} = 0$.

Question 7:

Let the vectors $\vec{a}, \vec{b}, \vec{c}$ given as $\hat{a_1i} + \hat{a_2j} + \hat{a_3k}$, $\hat{b_1i} + \hat{b_2j} + \hat{b_3k}$, $\hat{c_1i} + \hat{c_2j} + \hat{c_3k}$. Then show that $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$

Solution:

We have

$$\vec{a} = \vec{a}_1 \vec{i} + \vec{a}_2 \vec{j} + \vec{a}_3 \vec{k}$$

$$\vec{b} = \vec{b}_1 \vec{i} + \vec{b}_2 \vec{j} + \vec{b}_3 \vec{k}$$

$$\vec{c} = \vec{c}_1 \vec{i} + \vec{c}_2 \vec{j} + \vec{c}_3 \vec{k}$$

Then,

$$(\vec{b}+\vec{c}) = (b_1 + c_1)\hat{i} + (b_2 + c_2)\hat{j} + (b_3 + c_3)\hat{k}$$

Now,

$$\vec{a} \times (\vec{b} + \vec{c}) \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 + c_1 & b_2 + c_2 & b_3 + c_3 \end{vmatrix} = \begin{pmatrix} i [a_2(b_3 + c_3) - a_3(b_2 + c_2)] - j [a_1(b_3 + c_3) - a_3(b_1 + c_1)] \\ + k [a_1(b_2 + c_2) - a_2(b_1 + c_1)] \end{pmatrix}$$
$$= \begin{pmatrix} i [a_2b_3 + a_2c_3 - a_3b_2 - a_3c_2] + j [-a_1b_3 - a_1c_3 + a_3b_1 + a_3c_1] \\ + k [a_1b_2 + a_1c_2 - a_2b_1 - a_2c_1] \end{pmatrix}$$

Also,

$$\overrightarrow{a} \times \overrightarrow{b} = \begin{bmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix}$$

$$= i[a_2b_3 - a_3b_2] + j[a_3b_1 - a_1b_3] + k[a_2b_2 - a_2b_1]$$

$$\overrightarrow{a} \times \overrightarrow{c} = \begin{bmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \end{bmatrix}$$

$$= i[a_2c_3 - a_3c_2] + j[a_3c_1 - a_1c_3] + k[a_2c_2 - a_2c_1]$$

Therefore,

$$(\vec{a} \times \vec{b}) + (\vec{a} \times \vec{c}) = \begin{pmatrix} i[a_2b_3 - a_3b_2] + j[a_3b_1 - a_1b_3] + k[a_2b_2 - a_2b_1] \\ + i[a_2c_3 - a_3c_2] + j[a_3c_1 - a_1c_3] + k[a_2c_2 - a_2c_1] \end{pmatrix}$$

$$= \begin{pmatrix} i[a_2b_3 + a_2c_3 - a_3b_2 - a_3c_2] + j[-a_1b_3 - a_1c_3 + a_3b_1 + a_3c_1] \\ + k[a_1b_2 + a_1c_2 - a_2b_1 - a_2c_1] \end{pmatrix}$$

Thus,

$$\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$$

Hence proved.

Question 8:

If either a=0 or b=0, then $a \times b = 0$. Is the converse true? Justify your answer with an example.

Solution:

Let $\vec{a} = \hat{2}i + \hat{3}j + \hat{4}k$ and $\vec{b} = \hat{4}i + \hat{6}j + \hat{8}k$

Therefore,

$$\vec{a} \times \vec{b} = \begin{vmatrix} i & j & k \\ 2 & 3 & 4 \\ 4 & 6 & 8 \end{vmatrix}$$
$$= i(24 - 24) - j(16 - 16) + k(12 - 12)$$
$$= 0$$

Now,

$$\vec{a} = \sqrt{2^2 + 3^2 + 4^2} = \sqrt{29}$$

Thus,

Also,

$$\left| \vec{b} \right| = \sqrt{4^2 + 6^2 + 8^2} = \sqrt{116}$$

Thus,

 $\dot{b \neq 0}$

 $a \neq 0$

Hence, converse of the statement need not to be true.

Question 9:

Find the area of triangle with vertices A(1,1,2) B(2,3,5) and C(1,5,5).

Solution:

Vertices of the triangle are A(1,1,2) B(2,3,5) and C(1,5,5)

Hence,

$$\widehat{AB} = (2-1)\hat{i} + (3-1)\hat{j} + (5-2)\hat{k}$$

= $\hat{i} + 2\hat{j} + 3\hat{k}$
BC = $(1-2)\hat{i} + (5-3)\hat{j} + (5-5)\hat{k}$
= $\hat{-i} + 2\hat{j}$

Therefore,

Area of the triangle $ar(\Delta ABC) = \frac{1}{2} | \overrightarrow{AB} \times BC |$

Now,

$$\begin{array}{c} \blacksquare & \blacksquare & \blacksquare \\ AB \times BC = \begin{vmatrix} i & j & k \\ 1 & 2 & 3 \\ -1 & 2 & 0 \end{vmatrix} \\ = i(-6) - j(3) + k(2+2) \\ = -6i - 3j + 4k \\ \blacksquare \\ AB \times BC = \sqrt{(-6)^2 + (-3)^2 + 4^2} \\ = \sqrt{36 + 9 + 16} \\ = \sqrt{61} \end{array}$$

Therefore,

$$ar(\Delta ABC) = \frac{1}{2}\sqrt{61}$$
$$= \frac{\sqrt{61}}{2}$$

Question 10:

Find the area of the parallelogram whose adjacent sides are determined by the vector $\vec{a} = i - j + 3k$ and $\vec{b} = 2i - 7j + k$.

Solution: We have $\vec{a} = i - j + 3k$ and $\vec{b} = 2i - 7j + k$

Hence,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 3 \\ 2 & -7 & 1 \end{vmatrix}$$
$$= \hat{i}(-1+21) - \hat{j}(1-6) + \hat{k}(-7+2)$$
$$= 2\hat{0}\hat{i} + \hat{5}\hat{j} - \hat{5}\hat{k}$$
$$|\vec{a} \times \vec{b}| = \sqrt{20^2 + 5^2 + 5^2}$$
$$= \sqrt{400 + 25 + 25}$$
$$= 15\sqrt{2}$$

Thus, the area of parallelogram is $15\sqrt{2}$ square units.

Question 11:

Let the vectors \vec{a} and \vec{b} be such that $|\vec{a}| = 3$ and $|\vec{b}| = \frac{\sqrt{2}}{3}$, then $\vec{a \times b}$ is a unit vector, if the angle between \vec{a} and \vec{b} is

(A)
$$\frac{\pi}{6}$$
 (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$

Solution:

We have $|\vec{a}| = 3$, $|\vec{b}| = \frac{\sqrt{2}}{3}$ and $|\vec{a} \times \vec{b}| = 1$ Therefore,

$$\Rightarrow \left\| \overrightarrow{d} \right\| \overrightarrow{b} \right| \sin \theta = 1$$
$$\Rightarrow 3 \times \frac{\sqrt{2}}{3} \times \sin \theta = 1$$
$$\Rightarrow \sin \theta = \frac{1}{\sqrt{2}}$$
$$\Rightarrow \theta = \frac{\pi}{4}$$

Question 12:

Area of the rectangle having vertices A, B, C and D with position vectors $\hat{-i} + \frac{h}{2}j + \hat{4}k$ $i + \frac{h}{2}j + \hat{4}k$, $i - \frac{h}{2}j + \hat{4}k$ and $\hat{-i} - \frac{h}{2}j + \hat{4}k$, respectively is (A) $\frac{1}{2}$ (B) 1 (C) 2 (D) 4

Solution:

We have vertices
$$A\left(-i+\frac{r}{2}j+\hat{4}k\right)$$
, $B\left(i+\frac{r}{2}j+\hat{4}k\right)$, $C\left(i-\frac{r}{2}j+\hat{4}k\right)$ and $D\left(-i-\frac{r}{2}j+\hat{4}k\right)$.

Therefore,

$$\begin{array}{c} \blacksquare \\ AB = (1+1)\hat{i} + \left(\frac{1}{2} - \frac{1}{2}\right)j + (4-4)\hat{k} = \hat{2}i \\ \blacksquare \\ BC = (1-1)\hat{i} + \left(-\frac{1}{2} - \frac{1}{2}\right)j + (4-4)\hat{k} = -j
\end{array}$$

Now,

$$\overrightarrow{AB} \times \overrightarrow{BC} = \begin{cases} i & j & k \\ 2 & 0 & 0 \\ 0 & -1 & 0 \end{cases}$$
$$= \hat{k}(-2)$$
$$= -\hat{2}k$$
$$\overrightarrow{AB} \times \overrightarrow{BC} = \sqrt{(-2)^2}$$
$$= 2$$

So, area of the rectangle is 2 square units.

MISCELLANEOUS EXERCISE

Question 1:

Write down a unit vector in XY-plane, making an angle of 30° with the positive direction *x*-axis.

Solution:

Unit vector is $\vec{r} = \cos \hat{\theta} i + \sin \hat{\theta} j$, where θ is angle with positive x-axis. Therefore,

$$\vec{r} = \cos 30^{\circ} i + \sin 30^{\circ} j$$
$$= \frac{\sqrt{3}}{2} i + \frac{1}{2} j$$

Question 2:

Find the scalar components and magnitude of the vector joining the points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$.

Solution:

We have $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$

Therefore,

$$\begin{array}{c} PQ = (x_2 - x_1)i + (y_2 - y_1)j + (z_2 - z_1)k \\ PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2} \end{array}$$

Hence, the scalar components of the vector is $\{(x_2 - x_1) + (y_2 - y_1) + (z_2 - z_1)\}$ and magnitude of the vector is $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$.

Question 3:

A girl walks 4 km towards west, then she walks 3 km in a direction 30° east of north and stops. Determine the girl's displacement from her initial point of departure.

Solution:

Let O and B be the initial and final positions of the girl, respectively. Then, the girl's position can be shown by the below diagram:



We have:

$$\begin{aligned} \overrightarrow{DA} &= -4i \\ \overrightarrow{AB} &= i \left| \overrightarrow{AB} \right| \cos 60^\circ + j \left| \overrightarrow{AB} \right| \sin 60^\circ \\ &= i3 \times \frac{1}{2} + j3 \times \frac{\sqrt{3}}{2} \\ &= \frac{3}{2}i + \frac{3\sqrt{3}}{2}j \end{aligned}$$

By the triangle law of addition for vector,

$$\overrightarrow{OB} = \overrightarrow{OA} + \overrightarrow{AB}$$
$$= \left(-4i\right) + \left(\frac{3}{2}i + \frac{3\sqrt{3}}{2}j\right)$$
$$= \left(-4 + \frac{3}{2}\right)i + \frac{3\sqrt{3}}{2}j$$
$$= \left(\frac{-8+3}{2}\right)i + \frac{3\sqrt{3}}{2}j$$
$$= \frac{-5}{2}i + \frac{3\sqrt{3}}{2}j$$

Hence, the girl's displacement from her initial point of departure is $\frac{-5}{2}i + \frac{3\sqrt{3}}{2}j$.

Question 4:

If $\vec{a} = \vec{b} + \vec{c}$, then is it true that $|\vec{a}| = |\vec{b}| + |\vec{c}|$? Justify your answer.

Solution:

In $\triangle ABC$

$$CB = a, CA = b$$
 and $AB = c$



By triangle law of addition for vectors

$$a = b + c$$

By triangle inequality law of lengths

$\left| \overrightarrow{a} \right| < \left| \overrightarrow{b} \right| + \left| \overrightarrow{c} \right|$

Hence, it is not true that $\left| \vec{a} \right| = \left| \vec{b} \right| + \left| \vec{c} \right|$

Question 5:

Find the value of x for which x(i+j+k) is a unit vector.

Solution:

We have a unit vector $\hat{x(i+j+k)}$ Therefore,

$$\Rightarrow \left| x \left(\hat{i} + \hat{j} + k \right) \right| = 1$$

$$\Rightarrow \sqrt{x^2 + x^2 + x^2} = 1$$

$$\Rightarrow \sqrt{3x^2} = 1$$

$$\Rightarrow \sqrt{3x} = 1$$

$$\Rightarrow x = \pm \frac{1}{\sqrt{3}}$$

Question 6:

Find a vector of magnitude 5 units, and parallel to the resultant of the vectors $\vec{a} = 2i + 3j - k$ and $\vec{b} = i - 2j + k$.

Solution:

We have $\vec{a} = \hat{2}i + \hat{3}j - k$ and $\vec{b} = i - \hat{2}j + k$

Hence,

$$c = a + b$$

= $(2 + 1)i + (3 - 2)j + (-1 + 1)k$
= $3i + j$
 $|\vec{c}| = \sqrt{3^2 + 1^2}$
= $\sqrt{9 + 1}$
= $\sqrt{10}$

Therefore,

$$c = \frac{\overrightarrow{c}}{|c|} = \frac{\left(\widehat{3}i + j\right)}{\sqrt{10}}$$

So, a vector of magnitude 5 and parallel to the resultant of a and b is

$$\pm (c) = \pm 5 \left(\frac{1}{\sqrt{10}} (3i + j) \right)$$
$$= \pm \frac{3\sqrt{10}}{2} i \pm \frac{\sqrt{10}}{2} j$$

Question 7:

If $\vec{a} = i + j + k$, $\vec{b} = 2i - j + 3k$ and $\vec{c} = i - 2j + k$ find a unit vector parallel to the vector $\vec{2a} - \vec{b} + 3c$.

Solution:

We have $\vec{a} = i + j + k$, $\vec{b} = 2i - j + 3k$ and $\vec{c} = i - 2j + k$

Therefore,

$$\vec{2a-b+3c} = 2(\hat{i}+\hat{j}+\hat{k}) - (\hat{2}\hat{i}-\hat{j}+\hat{3}\hat{k}) + 3(\hat{i}-2\hat{j}+\hat{k})$$

= $\hat{2}\hat{i}+\hat{2}\hat{j}+\hat{2}\hat{k}-\hat{2}\hat{i}+\hat{j}-\hat{3}\hat{k}+\hat{3}\hat{i}-\hat{6}\hat{j}+\hat{3}\hat{k}$
= $\hat{3}\hat{i}-\hat{3}\hat{j}+\hat{2}\hat{k}$
 $|2\vec{a}-\vec{b}+3\vec{c}| = \sqrt{3^2+(-3)^2+2^2}$
= $\sqrt{9+9+4}$
= $\sqrt{22}$

So, the required unit vector is

$$\frac{2\overrightarrow{a-b+3c}}{|2a-b+3c|} = \frac{3i-3j+2k}{\sqrt{22}}$$
$$= \frac{3}{\sqrt{22}}i - \frac{3}{\sqrt{22}}j + \frac{2}{\sqrt{22}}k$$

Question 8:

Show that the points A(1,-2,-8), B(5,0,-2) and C(11,3,7) are collinear and find the ratio in which B divides AC.

Solution:

We have points A(1,-2,-8), B(5,0,-2) and C(11,3,7)

Therefore,

$$\begin{array}{l} \overrightarrow{AB} = (5-1)\hat{i} + (0+2)\hat{j} + (-2+8)\hat{k} = \hat{4}\hat{i} + \hat{2}\hat{j} + \hat{6}\hat{k} \\ \overrightarrow{BC} = (11-5)\hat{i} + (3-0)\hat{j} + (7+2)\hat{k} = \hat{6}\hat{i} + \hat{3}\hat{j} + \hat{9}\hat{k} \\ \overrightarrow{AC} = (11-1)\hat{i} + (3+2)\hat{j} + (7+8)\hat{k} = 10\hat{i} + \hat{5}\hat{j} + 1\hat{5}\hat{k} \\ \overrightarrow{AB} = \sqrt{4^2 + 2^2 + 6^2} = \sqrt{16 + 4 + 36} = \sqrt{56} = 2\sqrt{14} \\ \overrightarrow{BC} = \sqrt{6^2 + 3^2 + 9^2} = \sqrt{36 + 9 + 81} = \sqrt{126} = 3\sqrt{14} \\ \overrightarrow{AC} = \sqrt{10^2 + 5^2 + 15^2} = \sqrt{100 + 25 + 225} = \sqrt{350} = 5\sqrt{14} \end{array}$$

Now,

$$\begin{vmatrix} \Box B \\ AB \end{vmatrix} + \begin{vmatrix} BC \\ BC \end{vmatrix} = 2\sqrt{14} + 3\sqrt{14}$$
$$= 5\sqrt{14}$$
$$= \begin{vmatrix} \Box B \\ AC \end{vmatrix}$$

Thus, the points are collinear.

Let B divides AC in the ratio λ :1

Therefore,

$$\begin{split} & \bigoplus_{OB} = \frac{\lambda \overrightarrow{OC} + \overrightarrow{OA}}{(\lambda + 1)} \\ & \Rightarrow \hat{5}i - \hat{2}k = \frac{\lambda \left(1\hat{1}i + \hat{3}j + \hat{7}k\right) + \left(i - \hat{2}j - \hat{8}k\right)}{\lambda + 1} \\ & \Rightarrow (\lambda + 1) \left(\hat{5}i - \hat{2}k\right) = 11\hat{\lambda}i + 3\hat{\lambda}j + 7\hat{\lambda}k + i - \hat{2}j - \hat{8}k \\ & \Rightarrow 5(\lambda + 1)\hat{i} - 2(\lambda + 1)\hat{k} = (11\lambda + 1)\hat{i} + (3\lambda - 2)\hat{j} + (7\lambda - 8)\hat{k} \end{split}$$

On equating the corresponding components, we get

$$\Rightarrow 5(\lambda + 1) = (11\lambda + 1)$$
$$\Rightarrow 5\lambda + 5 = 11\lambda + 1$$
$$\Rightarrow 6\lambda = 4$$
$$\Rightarrow \lambda = \frac{2}{3}$$

Thus, the ratio is 2:3.

Question 9:

Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are (2a+b) and (a-3b) externally in the ratio 1:2. Also show that P is midpoint of the line segment RQ.

Solution:

We have $\overrightarrow{OP} = 2a + b$, $\overrightarrow{OQ} = a - 3b$

It is given that point R divides a line segment joining two points P and Q externally in the ratio 1: 2.

Then, on using the section formula, we get:

$$\overrightarrow{OR} = \frac{2(2a+b)-(a-2b)}{\overrightarrow{2-1}} = \frac{4a+2b-a-3b}{\overrightarrow{2-1}} = 3a+5b$$

Hence, the position vector of R is 3a + 5b

Thus, the position vector of midpoint of
$$RQ = \frac{\overrightarrow{OQ} + \overrightarrow{OR}}{2}$$

$$\frac{\overrightarrow{OQ} + \overrightarrow{OR}}{2} = \frac{(\overrightarrow{a} - 3\overrightarrow{b}) + (3\overrightarrow{a} + 5\overrightarrow{b})}{2}$$
$$= 2\overrightarrow{a} + \overrightarrow{b}$$
$$= \overrightarrow{OP}$$

Thus, P is the midpoint of line segment RQ.

Question 10:

The two adjacent sides of a parallelogram are $\hat{2}i - \hat{4}j + \hat{5}k$ and $i - \hat{2}j - \hat{3}k$. Find the unit vector parallel to its diagonal. Also, find its area.

Solution:

Diagonal of a parallelogram is a+b

$$\vec{a} + \vec{b} = (2+1)\hat{i} + (-4-2)\hat{j} + (5-3)\hat{k}$$
$$= \hat{3}\hat{i} - \hat{6}\hat{j} + \hat{2}\hat{k}$$

So, the unit vector parallel to the diagonal is

$$\vec{a+b} = \frac{\hat{3}i - \hat{6}j + \hat{2}k}{\sqrt{3^2 + (-6)^2 + 2^2}} = \frac{\hat{3}i - \hat{6}j + \hat{2}k}{\sqrt{3^2 + (-6)^2 + 2^2}} = \frac{\hat{3}i - \hat{6}j + \hat{2}k}{\sqrt{9 + 36 + 4}} = \frac{\hat{3}i - \hat{6}j + \hat{2}k}{7} = \frac{1}{7} (\hat{3}i - \hat{6}j + \hat{2}k)$$

Area of the parallelogram is $|\vec{a} + \vec{b}|$ Now,

$$\vec{a} \times \vec{b} = \begin{vmatrix} i & j & k \\ 2 & -4 & 5 \\ 1 & -2 & -3 \end{vmatrix}$$
$$= i(12+10) - j(-6-5) + k(-4+4)$$
$$= 2\hat{2}i + 1\hat{1}j$$
$$= 11(\hat{2}i + \hat{j})$$
$$|\vec{a} + \vec{b}| = 11\sqrt{2^2 + 1^2} = 11\sqrt{5}$$

So, area of parallelogram is $11\sqrt{5}$ square units.

Question 11:

Show that the direction cosines of a vector equally inclined to the axis OX, OY and OZ are

$$\pm \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$$

Solution:

Let a vector be equally inclined to OX, OY and OZ at an angle α

So, the DCs of the vectors are $\cos \alpha$, $\cos \alpha$ and $\cos \alpha$.

Therefore,

$$\cos^{2} \alpha + \cos^{2} \alpha + \cos^{2} \alpha = 1$$

$$\Rightarrow 3\cos^{2} \alpha = 1$$

$$\Rightarrow \cos^{2} \alpha = \frac{1}{3}$$

$$\Rightarrow \cos \alpha = \pm \frac{1}{\sqrt{3}}$$

$$\pm \begin{pmatrix} 1 & 1 & 1 \end{pmatrix}$$

Thus, the DCs of the vector are $\pm \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$.

Question 12:

Let $\vec{a} = i + \hat{4}j + \hat{2}k$, $\vec{b} = \hat{3}i - \hat{2}j + \hat{7}k$ and $\vec{c} = \hat{2}i - j + \hat{4}k$. Find a vector \vec{d} which is perpendicular to both \vec{a} and \vec{b} , and $\vec{c} \cdot \vec{d} = 15$.

Solution:

Let $\vec{d} = \hat{d_1}i + \hat{d_2}j + \hat{d_3}k$

Since, \vec{d} is perpendicular to both \vec{a} and \vec{b} , we have

 $\Rightarrow d_1 + 4d_2 + 2d_3 = 0 \qquad \dots (1)$

And

 $\vec{d}.\vec{a} = 0$

$$\overrightarrow{d.b} = 0$$

$$\Rightarrow 3d_1 - 2d_2 + 7d_3 = 0 \qquad \dots (2)$$

Also, it is given that

$$\overrightarrow{c.d} = 15$$

$$\Rightarrow 2d_1 - d_2 + 4d_3 = 15 \qquad \dots(3)$$

On solving equations (1),(2) and (3), we get

$$d_1 = \frac{160}{3}, d_2 = -\frac{5}{3}, d_3 = -\frac{70}{3}$$

Therefore,

$$\overrightarrow{d} = \frac{160}{3}i - \frac{5}{3}j - \frac{70}{3}k = \frac{1}{3} \left(160i - 5j - 70k \right)$$

Question 13:

The scalar product of the vector $\hat{i} + \hat{j} + \hat{k}$ with a unit vector along the sum of vectors $\hat{2}i + \hat{4}j - \hat{5}k$ and $\hat{\lambda}i + \hat{2}j + \hat{3}k$ is equal to one. Find the value of λ .

Solution:

$$\left(\hat{2}i+\hat{4}j-\hat{5}k\right)+\left(\hat{\lambda}i+\hat{2}j+\hat{3}k\right)=\left(2+\lambda\right)i+\hat{6}j-\hat{2}k$$

Therefore, unit vector along $(\hat{2}i+\hat{4}j-\hat{5}k)+(\hat{\lambda}i+\hat{2}j+\hat{3}k)$ is $\frac{(2+\lambda)\hat{i}+\hat{6}j-\hat{2}k}{\sqrt{(2+\lambda)^2+6^2+(-2)^2}} = \frac{(2+\lambda)\hat{i}+\hat{6}j-\hat{2}k}{\sqrt{4+4\lambda+\lambda^2+36+4}} = \frac{(2+\lambda)\hat{i}+\hat{6}j-\hat{2}k}{\sqrt{\lambda^2+4\lambda+44}}$

Scalar product of i+j+k with this unit vector is 1.

$$\Rightarrow \hat{i} + \hat{j} + \hat{k} \cdot \hat{k}$$

Question 14:

If a, b, c are mutually perpendicular vectors of equal magnitudes, show that the vector a+b+c is inclined to a, b, c.

Solution:

Since, $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular vectors of equal magnitudes Therefore, $\vec{a}, \vec{b} = \vec{b}, \vec{c} = \vec{c}, \vec{a} = 0$

And

$$\left| \overrightarrow{a} \right| = \left| \overrightarrow{b} \right| = \left| \overrightarrow{c} \right|$$

Let a+b+c be inclined to a,b,c at angles $\theta_1,\theta_2,\theta_3$ respectively.

$$\cos\theta_{1} = \frac{\overrightarrow{(a+b+c)}, \overrightarrow{a}}{|a+b+c||a|} = \frac{\overrightarrow{(a+b+c)}, \overrightarrow{a}}{|a+b+c||a|} = \frac{\overrightarrow{(a+b+c)}}{|a+b+c||a|} = \frac{\overrightarrow{(a+$$

Since $|\vec{a}| = |\vec{b}| = |\vec{c}|, \cos \theta_1 = \cos \theta_2 = \cos \theta_3$

Thus, $\theta_1 = \theta_2 = \theta_3$

Question 15: Prove that $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a}|^2 + |\vec{b}|^2$ if and only if \vec{a} and \vec{b} are perpendicular, given $\vec{a} \neq 0, \vec{b} \neq 0$.

Solution:

$$(a+b).(a+b) = |a|^2 + |b|^2$$

 $\Rightarrow a.a + a.b + b.a + b.b = |a|^2 + |b|^2$
 $\Rightarrow |a|^2 + 2a.b + |b|^2 = |a|^2 + |b|^2$
 $\Rightarrow 2a.b = 0$
 $\Rightarrow a.b = 0$

Thus, \vec{a} and \vec{b} are perpendicular.

Question 16:

If θ is the angle between two vectors \vec{a} and \vec{b} , then $\vec{a},\vec{b} \ge 0$ only when

(A) $0 < \theta < \frac{\pi}{2}$ (B) $0 \le \theta \le \frac{\pi}{2}$ (C) $0 < \theta < \pi$ (D) $0 \le \theta \le \pi$

Solution:

$$\vec{a}.\vec{b} \ge 0$$

$$\Rightarrow |\vec{a}||\vec{b}|\cos\theta \ge 0$$

$$\Rightarrow \cos\theta \ge 0$$

$$\Rightarrow 0 \le \theta \le \frac{\pi}{2}$$

$$[\because |\vec{a}| \ge 0 \text{ and } |\vec{b}| \ge 0]$$

Hence $a.b \ge 0$ if $0 \le \theta \le \frac{\pi}{2}$

Thus, the correct option is B.

Question 17:

Let \vec{a} and \vec{b} be two unit vectors and θ is the right angle between them. Then $\vec{a} + \vec{b}$ is a unit vector

(A)
$$\theta = \frac{\pi}{4}$$
 (B) $\theta = \frac{\pi}{3}$ (C) $\theta = \frac{\pi}{2}$ (D) $\theta = \frac{2\pi}{3}$

Solution:

We have \vec{a} and \vec{b} , two unit vectors and θ is the angle between them.

Then,

$$\left| \overrightarrow{a} \right| = \left| \overrightarrow{b} \right| = 1$$

Now, $\vec{a} + \vec{b}$ is a unit vector if $|\vec{a} + \vec{b}| = 1$ Therefore, $\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 1$ $\Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = 1$ $\Rightarrow |\vec{a}|^2 + 2\vec{a} \cdot \vec{b} + |\vec{b}|^2 = 1$ $\Rightarrow 1^2 + 2|\vec{a}||\vec{b}|\cos\theta + 1^2 = 1$ $\Rightarrow 1 + 2(1)(1)\cos\theta + 1 = 1$ $\Rightarrow \cos\theta = -\frac{1}{2}$ $\Rightarrow \theta = \frac{2\pi}{2}$

Hence, $\vec{a} + \vec{b}$ is a unit vector if $\theta = \frac{2\pi}{2}$

Thus, the correct option is D.

Question 18:

The value of $i.(j \times k) + j.(i \times k) + k(i \times j)$ is (A) 0 (B) -1 (C) 1 (D) 3

Solution:

$$\hat{i}.(\hat{j}\times\hat{k}) + \hat{j}.(\hat{i}\times\hat{k}) + \hat{k}(\hat{i}\times\hat{j}) = \hat{i}.\hat{i}+\hat{j}.(\hat{-j}) + \hat{k}.\hat{k}$$
$$= 1 - 1 + 1$$
$$= 1$$

Thus, the correct option is C.

Question 19:

If θ is the angle between any two vectors \vec{a} and \vec{b} , then $|\vec{a}\cdot\vec{b}| = |\vec{a}\times\vec{b}|$ when θ is equal to

(A) 0 (B)
$$\frac{\pi}{4}$$
 (C) $\frac{\pi}{2}$ (D) *n*

Solution:

Let θ be the angle between two vectors \vec{a} and \vec{b} .

Then, without loss of generality, \vec{a} and \vec{b} are non-zero vectors, so that $|\vec{a}|$ and $|\vec{b}|$ are positive.

Now,

$$\Rightarrow |\vec{a}.\vec{b}| = |\vec{a} \times \vec{b}|$$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta = |\vec{a}| |\vec{b}| \sin \theta$$

$$\Rightarrow \cos \theta = \sin \theta$$

$$\Rightarrow \tan \theta = 1$$

$$\Rightarrow \theta = \frac{\pi}{4}$$

So,
$$\left| \vec{a.b} \right| = \left| \vec{a \times b} \right|$$
 when $\theta = \frac{\pi}{4}$

Thus, the correct option is B.